LOUISIANA DEPARTMENT OF TRANSPORTATION AND DEVELOPMENT

PUBLIC WORKS AND WATER RESOURCES DIVISION

WATER RESOURCES SECTION

94° 00′

33° 00′

Potentiometric Surface--(Sheet 3 of 3)

Brantly, J.A., Seanor, R.C., and McCoy, K.L., 2002, Louisiana Ground-Water Map No. 13: Hydrogeology and potentiometric surface, October 1996, of the Sparta aquifer in northern Louisiana

below land surface datum

Land surface datum: 155 ft. sea level

Well: Ou-443

Well: Ou-444

Screen interval: 850-960 ft. below land surface datum

Land surface datum: 118 ft. sea level

Screen interval: 660-670 ft.,

below land surface datum

Land surface datum: 118 ft. sea level

IN FEET

ER

2000

-102

2000

POTENTIOMETRIC SURFACE

The generalized potentiometric surface of the Sparta aquifer, October 1996 (fig. 9), was constructed using water-level measurements from 55 wells (table 1) completed in the middle or lower part of the aquifer. The altitude of water-level contours ranged from about 220 ft above sea level in the outcrop area to about 220 ft below sea level in Ouachita Parish (fig. 9). Water levels are lowest (cones of depression) in areas affected by major pumping centers. Flow lines show ground-water movement is toward major pumping centers that include West Monroe (21 Mgal/d) and Hodge (10 Mgal/d) (fig. 9) in Louisiana. The Sparta aquifer also is affected by pumpage in the Ruston (6.9 Mgal/d), Minden (2.6 Mgal/d), and Winnfield (2.4 Mgal/d) areas of Louisiana, as well as pumpage near El Dorado, Arkansas (5.8 Mgal/d) (T.W. Holland, U.S. Geological Survey, oral commun., 2002). Water-level data collected subsequent to the data used for this potentiometric-surface map indicate the presence of a cone of depression in the Ruston area.

In the outcrop area in Webster Parish, well Wb-399 (fig. 10) shows only seasonal fluctuations in water levels for the period of record from 1979 to 2000 and no net increase or decrease. This well is typical of wells located in the outcrop area of the Sparta aquifer. Well Wb-399 is screened between a depth of 288 and 298 ft.

Declines in water levels in the Sparta aquifer are shown at well Cl-149 (fig. 11), located in Claiborne Parish near the Louisiana-Arkansas State line, and well L-26 (fig. 12), located north of Ruston in Lincoln Parish. For the periods of record, water levels in wells Cl-149 and L-26 have declined approximately 1.5 and 2.0 ft/yr, respectively. These declines are a result of long-term withdrawals from the aquifer.

Regional decline of water levels in the Sparta aquifer in Ouachita Parish and adjacent parishes occurs due to pumpage in West Monroe. Wells Ou-443, 444, and 445, located near West Monroe, are completed at different depths and monitor water levels in the West Monroe and surrounding areas. All three wells have shown an average water-level decline of about 2.0 ft/yr for the period of record shown. Hydrographs for these wells indicate water levels undergo very similar fluctuations (fig. 13). Water levels in these wells indicate that hydraulic connection exists among sands in the Sparta aquifer.

SUMMARY AND CONCLUSIONS

The Sparta aquifer is the principal source of ground water for Bienville, Claiborne, Jackson, Lincoln, Morehouse, Ouachita, Union, Webster, and Winn Parishes. Water levels throughout much of the aquifer have declined steadily since the early 1920's when water withdrawals for industrial purposes began. In 2000, the aquifer provided a total of 68 million gallons per day for public supply, industry, power generation, rural domestic supply, livestock, irrigation, and aquaculture. From 1990 to 2000, withdrawals from the aquifer increased 6 percent.

The Sparta aquifer framework consists of imperfectly connected sands with interbedded clays which may act as separate hydraulic units for short periods of time, but due to the depositional nature of the aquifer, act as one unit regionally over longer periods of time. Pumping from the massive sands in the lower part of the aquifer may affect water levels in the upper sands.

For most of the Sparta aquifer, the altitude of the top ranges from approximately 200 feet above sea level near the outcrop area to approximately 400 feet below sea level near the downdip limit of freshwater, and the top of the aquifer generally dips east and southeast at a rate of 5 to 25 feet per mile. The level in the outcrop area to approximately 1,000 feet below sea level near the downdip limit of freshwater, and the base of the aquifer dips east and southeast at a rate of 25 to 50 feet per mile.

Five hydrogeologic sections, oriented west-to-east or north-to-south, show the spatial extent of sands within the Sparta aquifer. Along the sections, the aquifer ranges in thickness from approximately 200 feet adjacent to the outcrop areas to approximately 725 feet near the downdip limit of freshwater.

The potentiometric surface in October 1996 shows that water levels in the Sparta aquifer generally have declined, except in the outcrop area where water levels have remained relatively constant. Altitude of water levels, which range from about 220 feet above sea level to about 220 feet below sea level, are highest in the western outcrop area and lowest in areas affected by major pumping centers.

Table 1. Water level data used to construct the notantiametric surface of

Well number	Well depth (feet below land surface)	Water level (feet below land surface)	Water level (feet above or below sea level)	Date measured
		Bienville	Parish	
Bi-76	110	50.46	229.54	10-01-96
Bi-100	90	33.21	186.79	10-16-96
Bi-112	348	115.82	99.18	10-01-96
Bi-144	630 472	236.42	83.58	10-01-96
Bi-166 Bi-186	216	181.43 41.97	78.57 138.03	10-01-96 10-16-96
Bi-192	153	74.49	210.51	10-16-96
Bi-216	300	16.70	183.30	10-01-96
		Caldwell	Parish	
Ca-86B	545	79.28	80.72	10-01-96
		Claiborne	Parish	
C1-9	670	270.01	89.99	10-09-96
Cl-58 Cl-111	488 570	140.01 255.46	109.99 44.54	10-09-96 10-08-96
Cl-1116	684	259.73	-14.73	10-10-96
Cl-148	625	196.80	-6.80	10-10-96
Cl-149	736	294.36	-64.36	10-10-96
		Jackson l	Parish	
Ja-49	570	201.19	-41.19	10-02-96
Ja-147 Ja-148	703 578	246.47 225.15	-26.47 19.85	10-16-96 10-02-96
		Lincoln I	Parish	
L-26	686	180.27	-25.27	10-10-96
L-68	770	249.60	-69.60	10-10-96
L-113	750	317.56	37.44	10-02-96
L-117	551	Morehouse	-51.85	10-15-96
M- 5	960	и		10.04.04
Mo-5 Mo-342	860 620	153.52 89.85	-36.08 -1.51	10-04-96 10-03-96
Mo-350	740	105.56	6.52	10-03-96
		Ouachita	Parish	
Ou-80	176	283.76	-223.76	10-10-96
Ou-401 A	A 397 750	106.13 58.98	-43.85 4.32	10-10-96 10-10-96
Ou-404	167	80.32	-19.43	10-15-96
Ou-406	681	157.25	-90.50	10-10-96
Ou-444	670	229.09	-111.09	10-23-96
Ou-488	255	332.73	-52.73	10-23-96
II. 26	745	Union Pa		10.04.00
Un-26 Un-78	745 727	184.00 208.60	-50.08 -33.60	10-04-96 10-24-96
Un-79	749	153.40	-35.40	10-24-96
Un-83	326	174.00	-52.00	10-16-96
Un-84	696	248.70	-38.70	10-24-96
Un-86 Un-134	655 585	100.63 307.13	-10.63 -87.13	10-16-96 10-23-96
		Webster 1		
Wb-127	408	61.37	208.63	10-21-96
Wb-164	769	64.32	155.68	10-07-96
Wb-219	136	5.17	184.83	10-08-96
Wb-271 Wb-285	220 608	84.25	193.75 159.32	10-09-96
Wb-285 Wb-326	470	180.68 114.51	159.32	10-08-96 10-04-96
Wb-338	425	94.89	135.11	10-04-96
Wb-349	263	70.38	109.62	10-07-96
Wb-359	687	79.68	200.32	10-09-96
	298 544	45.24 17.20	159.76 142.80	10-04-96 10-09-96
Wb-399 Wb-415				
Wb-399		Winn Pa	nrish	
Wb-399 Wb-415 W-28	606	52.14	52.86	10-01-96
Wb-399 Wb-415 W-28 W-144B	550	52.14 37.55	52.86 102.45	10-01-96
Wb-399 Wb-415 W-28		52.14	52.86	10-01-96 10-01-96 10-01-96 10-01-96

SELECTED REFERENCES


- Clark, G.C., 1959, Interior salt domes of Texas, Louisiana, and Mississippi: Interior salt domes and Tertiary stratigraphy of Louisiana: Shreveport Geological Society, 1960, 147 p.
- Lovelace, J.K., 1991, Water use in Louisiana, 1990: Louisiana Department of Transportation and Development Water Resources Special Report no. 6, 131 p.
- Lovelace, J.K., and Johnson, P.M., 1996, Water use in Louisiana, 1995: Louisiana Department of Transportation and Development Water Resources Special Report no. 11, 127 p. McWreath, H. C. III, Nelson, J. D., and Fitzpatrick, D. J., 1991, Simulated response
- to pumping stresses in the Sparta aquifer, northern Louisiana and southern Arkansas: Louisiana Department of Transportation and Development Water Resources Technical Report no. 51, 51 p.
- Payne, J.N., 1968, Hydrologic significance of the lithofacies of the Sparta Sand in Arkansas, Louisiana, Mississippi, and Texas: U.S. Geological Survey Professional Paper 569-A, p. A1-A17.
- Page, L.V., and May, H.G., 1964, Water resources of Bossier and Caddo Parishes, Louisiana: Department of Conservation, Louisiana Geological Survey, and Louisiana Department of Public Works Water Resources Bulletin no. 5, 105 p.
- Ryals, G.N., 1980, Potentiometric surface maps of the Sparta Sand; northern Louisiana and southern Arkansas, 1900, 1965, 1975, and 1980: U.S. Geological Survey Open-File Report 80-1180, 1 sheet.
- ----, 1982, Ground-water resources of the Arcadia-Minden area, Louisiana: Louisiana Department of Transportation and Development, Office of Public Works Water Resources Technical Report no. 28, 35 p.
- Sanford, T.H., Jr., 1973a, Ground-water resources of Morehouse Parish, Louisiana: Department of Conservation, Louisiana Geological Survey, and Louisiana Department of Public Works Resources Bulletin no. 19, 90 p.

- ----1973b, Water resources of the Ruston area, Louisiana: Louisiana Department of Public Works Water Resources Technical Report no. 8, 32 p.
- Smoot, C.W., and Seanor, R.C., 1991, Louisiana ground-water map no. 3: Potentiometric surface, 1989, and water-level changes, 1980-89, of the Sparta aquifer in north-central Louisiana: U.S. Geological Survey Water-Resources Investigations Report 90-4183, 2 sheets.
- Snider, J.L., Calandro, A.J., and Shampine, W.J., 1972, Water resources of Union Parish, Louisiana: Department of Conservation, Louisiana Geological Survey, and Louisiana Department of Public Works Water Resources Bulletin no. 17, 68 p.
- Stuart, C.G., Knochenmus, Darwin, and McGee, B.D., 1994, Guide to Louisiana's ground-water resources: U.S. Geological Survey Water-Resources Investigations Report 94-4085, 55 p.
- Trudeau, D.A., Buono, Anthony, 1985, Projected effects of proposed increased pumpage on water levels and salinity in the Sparta aquifer near West Monroe, Louisiana: Louisiana Department of Transportation and Development Water Resources Technical Report no. 39, 70 p.
- U.S. Environmental Protection Agency, 1994, Drinking water regulations and health advisories: U.S. Environmental Protection Agency, Office of Water, May 1994, EPA 822-R-94-001,11 p.
- Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Louisiana Department of Transportation and Development - U.S. Geological Survey Water Resources Cooperative Program







2000

ARKANSAS

UNION

Figure 12. Water level in well L-26 (Lincoln Parish)


Well: Ou-445

Screen interval: 444-454 ft.,

below land surface dutum Land surface dutum: 118 ft. sea level

Figure 13. Water level in well Ou-443, Ou-444 and Ou-445 (Ouachita Parish).

93° 00′



Louisiana Ground-Water Map No. 13 Hydrogeology and Potentiometric Surface, October 1996, of the Sparta Aquifer in Northern Louisiana

Land surface datum: 205 ft. sea level

Figure 10. Water level in well Wb-399 (Webster Parish)

Jeffrey A. Brantly, Ronald C. Seanor, and Kaycee L. McCoy

2002